

# PARASTAR™

(fipronil)



Kills Fleas, Ticks & Chewing Lice.  
Fast-acting. Long-lasting. Waterproof.  
Contains fipronil, the active ingredient in Frontline® Top Spot.®\*

## KEEP OUT OF REACH OF CHILDREN CAUTION

ACTIVE INGREDIENTS:  
Fipronil.....9.7%  
INERT INGREDIENTS.....90.3%  
TOTAL.....100.0%

NET CONTENTS: Three applicators

Parastar™ contains fipronil to effectively kill fleas and ticks, including those that may transmit Lyme disease.




- Kills fleas • Kills deer ticks • Kills brown dog ticks
- Kills American dog ticks • Kills lone star ticks
- Kills chewing lice • Controls mites that may cause sarcoptic mange

### How to Open:

1. Remove product tubes from the package.
2. Separate one tube from the others. Holding the tube with notched end pointing up and away from the face and body, cut off the narrow end at the notches with scissors.

### How to Apply:

1. Invert tube over dog and use open end to part dog's hair.
2. Squeeze tube firmly to apply all of the solution to the dog's skin as a spot to the dog's back between the shoulder blades.



## READ ENTIRE LABEL BEFORE EACH USE. USE ONLY ON DOGS AND PUPPIES OVER 8 WEEKS OF AGE.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Do not allow children to apply product.** TO PREVENT HARM TO YOU AND YOUR DOG, READ ENTIRE LABEL BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY. FOR EXTERNAL USE ON DOGS ONLY. DO NOT USE ON RABBITS. DO NOT USE ON OTHER ANIMALS.

Do not use on puppies under 8 weeks of age. Weigh your dog to be sure you are applying the right dose formulated for the weight of your dog. Consult a veterinarian before using this product on debilitated, aged, or medicated dogs. Can be used on breeding, pregnant, or lactating dogs. Consult a veterinarian before using on dogs with known organ dysfunction. Dogs may experience some temporary irritation at the site of product application. Certain medications can interact with pesticides. Sensitivities may occur after using ANY pesticide product on dogs. If signs of sensitivity occur bathe your dog with a mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

**How to open:** Remove product tubes from the package. Separate one tube from the others. Holding the tube with notched end pointing up and away from the face and body, cut off the narrow end at the notches with scissors.

**How to apply:** While holding the dog with one hand, use the other hand to apply the solution. Invert the tube and use the open end to part the dog's hair. Apply contents of one tube of Parastar as a spot to the dog's back between the shoulder blades. Hold the dog for a few seconds to give the solution time to be absorbed into the dog's coat. When finished, wrap the tube in paper and put into the trash. Only one applicator per treatment is needed. Repeat application may be made if necessary, but do not apply more often than once every 30 days.

### FREQUENCY OF APPLICATION

Use Parastar monthly for control of flea, tick and chewing lice infestations. Studies show that Parastar kills fleas for up to three months. If your pet is at high risk for flea reinfestation, or in a highly infested environment, apply monthly. Apply monthly to control ticks and chewing lice. Parastar remains effective, after water immersion, or exposure to sunlight. Allow treated area to dry thoroughly. Do not reapply for 30 days.

**For Dogs Weighing up to 22 lbs.** Apply one tube 0.023 fl oz (0.67 mL) as a spot on the dog's back between the shoulder blades.

**For Dogs Weighing 23 to 44 lbs.** Apply one tube 0.045 fl oz (1.34 mL) as a spot on the dog's back between the shoulder blades.

**For Dogs Weighing 45 to 88 lbs.** Apply one tube 0.091 fl oz (2.68 mL) as a spot on the dog's back between the shoulder blades.

**For Dogs Weighing 89 to 132 lbs.** Apply one tube 0.136 fl oz (4.02 mL) as a spot on the dog's back between the shoulder blades.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal

**PESTICIDE STORAGE:** Do not remove tube from the pack until ready to use. Store in a cool (below 77°F) dry place inaccessible to children and pets. Do not refrigerate. Protect from direct sunlight.

**CONTAINER DISPOSAL:** Non-refillable container. Do not reuse or refill this container. If empty: Place in the trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

### FIRST AID

If swallowed	Call a poison control center or a doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If in eyes	Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin	Wash with plenty of soap and water. Get medical attention if irritation persists.

### HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For more information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), you may call 1-800-637-0281.

### PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS. CAUTION.** Harmful if swallowed. Causes eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

**HAZARDS TO DOMESTIC ANIMALS.** For external use only. Do not use on puppies under 8 weeks of age. Individual sensitivities, while rare, may occur after using any pesticide product. Dogs may experience some temporary irritation at the site of product application. If signs persist, or become more severe within a few days of application, consult a veterinarian immediately. If your dog has an unusual reaction to the initial application, consult a veterinarian before repeating treatment. Certain medications can interact with pesticides. Consult a veterinarian before using on medicated, debilitated or aged dogs.

### WARRANTY

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of NOVARTIS ANIMAL HEALTH US, Inc. (NAHUS) or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold NAHUS and Seller harmless for any claims relating to such factors.

NAHUS warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or NAHUS, and Buyer and User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NAHUS MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with State law, in no event shall NAHUS or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF NAHUS AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF NAHUS OR SELLER, THE REPLACEMENT OF THE PRODUCT.

For Technical Information, call: 1-800-637-0281

Made in USA

Manufactured for  
Novartis Animal Health US, Inc.  
3200 Northline Avenue, Suite 300  
Greensboro, NC 27408

EPA Registration No. 70585-12  
EPA Est. No. 2517-TN-1



© 2011 Novartis Animal Health US, Inc. Parastar is a trademark of Novartis AG.

\*Parastar™ is not manufactured by or distributed by Merial. Frontline® Top Spot® is a registered trademark of Merial.